

Project IP3

Improved Cassava for the Developing world



**A
N
N
U
A
L

R
E
P
O
R
T**

2004



CONTENTS

OUTPUT 1

Genetic base of cassava and related *Manihot* species evaluated and available for cassava improvement: higher nutritional quality

Variation of quality traits in cassava roots evaluated in landraces and improved clones (<i>Euphytica</i>).	1-1
Variation in crude protein content in cassava (<i>Manihot esculenta Crantz</i>) roots.	1-9
Reduction or delay of post-harvest physiological deterioration in high-carotene cassava roots.	1-12
Effect of processing on carotenes present in cassava roots.	1-17
Evaluation of different storage conditions of carotenes from fresh and processed cassava roots.	1-24
Measurement of genotype by environment effect on carotene, Fe and Zn contents in cassava roots.	1-25

OUTPUT 2

Genetic base of cassava and related *Manihot* species evaluated and available for cassava improvement: higher commercial value

A cassava-breeding scheme based on the production of doubled-haploids	2-5
Mutagenesis and the "TILLING" system.	2-7
Recurrent selection to increase and reduce amylose proportion in the root starch of cassava.	2-9
High-capacity starch quality laboratory	2-11

OUTPUT 3

Development of new genetic stocks and improved gene pools for their evaluation in key target environments

Selection of progenitors based on previous cycle results and information from other outputs (i.e., resistance/tolerance, root quality traits, etc.).	3-1
Establishment of crossing blocks and production of recombinant seed from previously established blocks.	3-4
Generation and distribution of advanced breeding materials for National Programs.	3-6
Selection of recombinant progenies for broad and specific adaptation within major agro-ecosystems.	3-8
Precision of selection in early stages of cassava genetic improvement. (Manuscript submitted to Crop Science)	3-13

OUTPUT 4

Development of genetic stocks and improved gene pools adapted to the sub-humid environments

Evaluations and selections in the sub-humid environment.	4-1
--	-----

OUTPUT 5

Development of genetic stocks and improved gene pools adapted to the sub-humid environments

Evaluations and selections in the Acid Soils Environment **5-1**

OUTPUT 6

Development of genetic stocks and improved gene pools adapted to the sub-humid environments

Evaluations and selections in the Mid-altitude Valleys **6-1**

OUTPUT 7

Development of genetic stocks and improved gene pools adapted to the sub-humid environments

Evaluations and selections in Córdoba and Sucre Departments **7-1**

Evaluations and selections in Middle-Magdalena River Region. **7-4**

Evaluations and selections in Tolima-Huila Departments Region. **7-6**

Evaluations and selections in the Highlands Region. **7-7**

OUTPUT 8

Collaboration with other institutions, scientific meetings, and publications

Support national programs that have traditionally collaborated with CIAT in the development and improvement of cassava. **8-1**

Development of collaborative projects with partners in Africa, Asia and Latin America and the Caribbean. **8-3**

The collaboration with Colombia. **8-5**

Scientific meetings and publications. **8-10**

OUTPUT 9

Activities related with the maintenance of the germplasm bank of cassava and other *Manihot* species. Basic genetic studies

Maintenance of *Manihot* germplasm bank in the field. **9-1**

Evaluation of *M. esculenta* and related species from the germplasm collection for useful traits, particularly for higher protein content in the roots. **9-4**

Evaluation of segregation of carotene content in self-pollinated progenies from selected clones. **9-5**

Evaluation of segregation of traits related to Post-Harvest Physiological Deterioration. **9-7**

Development of a quantitative genetics estimate of the standard error for test for epistasis. **9-8**

OUTPUT 10

Breeding for insect and other arthropods resistance and development of alternative methods for their control

Evaluation of cassava germplasm for resistance to whiteflies (<i>Aleurotrachelus socialis</i>) during 2003-2004.	10-1
Evaluation of whitefly (<i>Aleurotrachelus socialis</i>) populations and damage on cassava genotypes from GM and CM families (developed for genome mapping studies for root dry matter) at Santander de Quilichao, 2004.	10-12
Intrinsic rate of increase of Biotype "B" <i>Bemisia tabaci</i> on two African cassava genotypes MNg 2 and MNg 11.	10-13
Studies on the biology and behavior of biotype "B" of <i>Bemisia tabaci</i> on a wild <i>Manihot</i> sp, <i>M. flabellifolia</i> .	10-19
Evaluation of cassava germplasm in several breeding and genetic trials for insect and mite pest damage at several localities on the Colombia Atlantic Coast.	10-23
Observations on the incidence, damage and behavior of whitefly (<i>A. socialis</i>), mites (<i>O. peruvianus</i>) and other arthropod pests in germplasm (breeding) trials on the Atlantic Coast of Colombia.	10-38
Cassava germplasm evaluations to identify resistance to the cassava green mite, <i>Mononychellus tanajoa</i> .	10-39
Testing of transgenic cassava (Africa genotype TMS 60444) plants displaying indications of resistance to the cassava hornworm, <i>Erinnyis ello</i> .	10-48
Determining the plant metabolites involved in whitefly (<i>Aleurotrachelus socialis</i>) resistant cassava varieties, MEcu 64, MEcu 72 and MPer 334.	10-51

OUTPUT 11

Disease Resistance in Cassava

Characterizing cassava genotypes for their reaction to super-elongation disease (SED) under greenhouse conditions, using different isolates	11-1
Characterizing cassava genotypes for their reaction to cassava bacterial blight (CBB) under greenhouse conditions, using different isolates	11-3
Evaluating cassava genotypes for their resistance to super-elongation disease (SED) in Santander de Quilichao, Department of Cauca, Colombia	11-6
Evaluating cassava genotypes growing in Pescador, Department of Cauca, Colombia, for their resistance to <i>Phytophthora</i> root rots (PRRs)	11-7
Evaluating of cassava genotypes growing in the departments of Sucre and Córdoba, for their resistance to FSD and SED.	11-8
Identifying the association between foliar resistance and root resistance to <i>Phytophthora tropicalis</i> . Determining resistance in roots and leaves to <i>P. tropicalis</i> during its penetration and post-penetration phases.	11-8
Determining the biochemical markers and agronomic traits associated with resistance to root rot caused by <i>Phytophthora tropicalis</i>	11-17
Determining the QTLs that most contribute to phenotypic variance of resistance to <i>Phytophthora</i> root rots, and identifying the linkage group(s) where they are located	11-36
Detecting phytoplasmas in cassava affected by frogskin disease (FSD), using nested PCR.	11-39

Identifying phytoplasmas by sequencing PCR products	11-44
Designing specific primers for high-specificity detection of a phytoplasma associated with frog skin disease (FSD) of cassava	11-48
Detecting phytoplasmas by electron microscopy	11-50
To develop and validate sustainable methods to prevent and control FSD and SED.	11-53
Multiplying cassava genotypes to ensure sufficient cuttings for disease-resistance evaluations	11-53
DNA sequence analysis of specific regions of cassava resistance genes analogs (RGAs).	11-53
Training researchers from Latin America, the Caribbean, and Africa on managing cassava diseases and research technology	11-55
Train students, farmers, technicians, and researchers through field days and meetings on modern, sustainable, cassava production systems in different regions of Colombia to manage major cassava diseases, emphasizing selection of stem cuttings	11-56
Publications in 2004	11-57
Two postgraduate theses in cassava for the Universidad Nacional de Colombia (Palmira) and the Universidad de los Andes (Bogotá, Colombia)	11-58
Two undergraduate theses currently being undertaken in cassava for the Universidad de Caldas, Manizales, Colombia	11-59
Concept notes and projects developed.	11-59

OUTPUT 12

Development and use of biotechnology tools for cassava improvement

Molecular Marker-Assisted Breeding for Resistance to the Cassava Mosaic Disease in Latin American Cassava Gene Pools	12-1
Molecular Marker-Assisted and Farmer Participatory Improvement of Cassava Germplasm for Farmer/Market Preferred Traits in Tanzania	12-3
Genetic Mapping of Genes Involved in the Biosynthesis of Beta-carotene	12-10
Progress in Genetic Mapping of Dry Matter Content (DMC) in Cassava	12-12
QTL Mapping of Cyanogenic Glucoside Content in a S ₁ Population derived from MTA18 and Candidate Gene Mapping of Two Cytochrome P-450 Biosynthetic Genes (CYP79D1 Y CYP79D2)	12-18
Development of Mapping Populations for Gene Tagging of Post Harvest Physiological Deterioration (PPD), Resistance to Hornworms and Whiteflies Found in Wild Relatives of Cassava	12-22
Generation Challenge Program: Comparison of Simple Sequence Repeats (SSR) and Diversity Array Technology (DArT) Markers for Structural Characterization of Diversity in Cassava	12-24
Generation Challenge Program: Assembling Germplasm and Molecular Markers Sets for Analysis of Structural Diversity in Cassava	12-28
Analysis of genetic diversity in a cassava germplasm collection from Cuba using SSR markers	12-34
Studies in Market Preferences of Cassava Cultivars in Malawi using SSR Markers	12-38
Data base of the Molecular Diversity Network of Cassava (MOLCAS)	12-41
Mining the Primary Gene Pool of Cassava: Introgression of High Root Protein from Accessions of <i>Manihot esculenta</i> sub spp <i>Fabellifolia</i> and <i>Manihot</i>	12-42

Tristis into Cassava	
Mining the Primary Gene Pool: Green Mites (CGM) Resistance Genes from <i>Manihot tristis</i> .	12-48
Identification of Naturally Occurring and Irradiation-Induced Mutant GBSSI Alleles of Cassava in a Heterozygous Genetic Background	12-50
Molecular Characterization of A Putative Waxy Cassava Starch GMO Obtained by Anti-sense Mediated Silencing of the Granule Bound Starch Synthase I (GBSSI) gene	12-55
Modification of Flowering in Cassava	12-58
Construction of a TME-3 Bacterial Artificial Chromosome (BAC) Library and Development of a BAC Contig around a CMD Resistance Gene	12-61
Isolation of Full-Length cDNA Clones of Transcripts Differentially Expressed Between Full-Sib Genotyped Resistant and Susceptible to the Cassava Mosaic Disease (CMD).	12-65
Embryo Rescue of Sexual Seeds from BC ₂ Families for Molecular Marker-Assisted Selection (MAS) for Resistance to Cassava Green Mites (CGM) and the Cassava Mosaic Disease (CMD)	12-68
Dissemination of Improved Cassava Varieties and Management of Genetic Stocks as Tissue Culture Plantlets	12-71
A Simple Method for the Rapid Multiplication of Clean Cassava Planting Material	12-74
Training	12-77
Trips	12-78
Publications	12-78
Project Funded and in Review with Donors	12-79

OUTPUT 13

[Integrated cassava-based cropping systems in Asia: Widespread adoption of farming practices that enhance sustainability](#)

Soil fertility maintenance through the application of chemical fertilizers, or the use of intercropping, green manuring, alley cropping and crop rotations.	13-1
Development of efficient and economical soil preparation practices.	13-4
Determination of the response to various methods of application of Zn in calcareous soils.	13-8
Evaluation of cassava varieties and determination of optimum plant spacing for cassava leaf production.	13-10
Conducting FPR trials on varieties, fertilization, weed control, green manures, intercropping, erosion control and pig feeding in Thailand, Vietnam and China.	13-14
Enhancing adoption of new varieties and improved management practices through farmer participatory research (FPR) and extension (FPE) activities.	13-21
Assessing the impact of the project on adoption of new technologies in Thailand and Vietnam	13-23
Exploring institutional arrangements for collaboration in the new Nippon Foundation-funded cassava project in Laos and Cambodia, and the ACIAR-funded cassava project in Indonesia and East Timor.	13-27
Implementing the new Nippon Foundation-funded Cassava Project in Lao PDR and Cambodia	13-29